

Interchange of filtered 2-colimits and finite 2-limits

Delphine Dupont

Abstract

In this paper we go into the study of 2-limit and 2-colimit in the 2-category $\mathcal{CAT}$ the category of small categories. In particular we show the commutation of filtered 2-colimits and finite 2-limits. It is a generalization of a classical result in category theory (see for example [1]).

1 Introduction

Recently the 2-category theory has developed rapidly. It gives a very useful language in many field of mathematics. One of the main notions is the 2-limits and 2-colimits. For example the stalk of a stack is a 2-colimit. In this paper we generalize a classical result in category theory.

In a first part we recall the definition of 2-limit and 2-colimit in $\mathcal{CAT}$, the 2-category of small categories. We give an explicit description of a filtered 2-colimit and of a 2-limit. But we do not express the morphisms of these categories in a classical way. We give them as elements of a colimit or a limit.

In a second part we prove the interchange of filtered 2-colimits and finite 2-limits. To prove this theorem we use the explicit expression of the categories $2 \lim_{i \in \mathcal{I}} 2 \lim_{j \in \mathcal{J}} \mathbf{a}(i, j)$ and $2 \lim_{j \in \mathcal{J}} 2 \lim_{i \in \mathcal{I}} \mathbf{a}(i, j)$.

We do not recall the definitions of a 2-category, a 2-functor (some times called pseudo functor in the literature), a 2-natural transformation of functors and a 2-modification. The reader can find them in the chapter 7 of [1], the appendix of the paper [3] and the paper [2].

Let us recall the definition of a filtered category :

Definition 1. *A category $\mathcal{I}$ is filtered if it satisfies the conditions (i)-(iii) below.*

- (i) $\mathcal{I}$ is non empty,
- (ii) for any i and j in $\mathcal{I}$, there exist $k \in \mathcal{I}$ and morphisms $i \rightarrow k, j \rightarrow k$,
- (iii) for any parallel morphisms $s, s' : i \rightrightarrows j$ there exists a morphism $h : j \rightarrow k$ such that $h \circ s = h \circ s'$.

2 2-colimits and 2-limits

Let us first recall the definitions of a 2-limit and a 2-colimit. This part is inspired by the appendix of [3]. We cite [2] as a classical reference.

Definition 2. Let $\mathcal{I}$ be a small category, and $\mathbf{b} : \mathcal{I} \rightarrow \mathcal{CAT}$ a 2-functor. The system $\mathbf{b}$ admits a 2-colimit if and only if there exist :

- a category $2\varinjlim_{i \in \mathcal{I}} \mathbf{b}(i)$ and
- a 2-natural transformation $\sigma : \mathbf{b} \rightarrow 2\varinjlim_{i \in \mathcal{I}} \mathbf{b}(i)$

such that for all category $\mathcal{C}$ the functor :

$$(\sigma \circ) : \underline{Hom}_{\mathcal{C}}(2\varinjlim_{i \in \mathcal{I}} \mathbf{b}(i), \mathcal{C}) \rightarrow \underline{Hom}(\mathbf{b}, \mathcal{C})$$

is an equivalence of categories.

We say that a 2-colimit has the strong factorisation property if $(\sigma \circ)$ is an isomorphism of categories.

More concretely, a 2-functor $\mathbf{b} : \mathcal{I} \rightarrow \mathcal{CAT}$ admits a 2-colimit if and only if there exist :

- a category $2\varinjlim_{i \in \mathcal{I}} \mathbf{b}(i)$,
- functors $\sigma_i : \mathbf{b}(i) \rightarrow 2\varinjlim_{i \in \mathcal{I}} \mathbf{b}(i)$ for any $i \in \mathcal{I}$,
- and a natural equivalence $\Theta_s^\sigma : \sigma_i \xrightarrow{\sim} \sigma_j \mathbf{b}(s)$ for any morphism $s : i \rightarrow j$ of $\mathcal{I}$ visualized by :

$$\begin{array}{ccc} & 2\varinjlim_{i \in \mathcal{I}} \mathbf{b}(i) & \\ \sigma_i \swarrow & & \searrow \sigma_j \\ \mathbf{b}(i) & \xrightarrow{\mathbf{b}(s)} & \mathbf{b}(j) \end{array}$$

$\sim$
 θ_s


such that $\sigma : \mathbf{b} \rightarrow 2\varinjlim_{i \in \mathcal{I}} \mathbf{b}(i)$ is a 2-natural transformation.

Moreover these data satisfy condition assuring that $(\sigma \circ)$ is an equivalence. The first one translates the fact that $(\sigma \circ)$ is essentially surjective and the second one that $(\sigma \circ)$ is fully faithful.

- For any category $\mathcal{C}$, any morphism of functors $\rho : \mathbf{b} \rightarrow \mathcal{C}$, there exist a functor :

$$F : 2\varinjlim_{i \in \mathcal{I}} \mathbf{b}(i) \longrightarrow \mathcal{C}$$

and an isomorphism $\varphi^F : \rho \rightarrow F\sigma$, which is a modification given, for all $i \in \mathcal{I}$, by a natural equivalence $\varphi_i^F : \rho_i \rightarrow F \circ \sigma_i$. This may be visualized by :



The compatibility condition is given by :

$$(F \bullet \theta_s^\sigma) \circ \varphi_i^F = (\varphi_j^F \bullet a(s)) \circ \theta_s^\rho$$

The pair (F, φ^F) is called a lax factorization of the system ρ .

- Let ρ and ρ' be two 2-natural transformations and $\lambda : \rho \rightarrow \rho'$ a modification. Then for any lax factorization $F : 2\varinjlim_{i \in \mathcal{I}} a(i) \rightarrow C$ of ρ and $G : 2\varinjlim_{i \in \mathcal{I}} b(i) \rightarrow C$ of ρ' there exists a unique natural transformation $\Lambda : F \rightarrow G$ such that :

$$\varphi_i^G \circ \lambda_i = (\Lambda_i \bullet \sigma_i) \circ \varphi_i^F$$

If the 2-colimit has the strong factorization property then there exists a unique factorization such that φ^F is the identity.

A contravariant 2-functor $\mathfrak{c} : \mathcal{I} \rightarrow \mathcal{CAT}$ is a 2-functor $\mathfrak{c} : \mathcal{I}^{op} \rightarrow \mathcal{CAT}$. A 2-limit is construction dual of the 2-colimit. Hence, we have the following definition :

Definition 3. Let $\mathcal{I}$ be a small category and $\mathfrak{c} : \mathcal{I} \rightarrow \mathcal{CAT}$ a contravariant 2-functor. The system $\mathfrak{c}$ admits a 2-limit if and only if there exist :

- a category $2\varprojlim_{i \in \mathcal{I}} \mathfrak{c}(i)$ and
- a 2-natural transformation $\sigma : 2\varprojlim_{i \in \mathcal{I}} \mathfrak{b}(i) \rightarrow \mathfrak{c}$

such that for all category $\mathcal{C}$ the functor :

$$(\circ \sigma) : \underline{Hom}(\mathfrak{c}, \mathcal{C}) \rightarrow \underline{Hom}_{\mathcal{C}}(2\varprojlim_{i \in \mathcal{I}} \mathfrak{b}(i), \mathcal{C})$$

is an equivalence of categories.

We say that a 2-limit has the strong factorisation property if $(\circ \sigma)$ is an isomorphism of categories.

It is well known that the 2-category $\mathcal{CAT}$ is complete and co-complete. The proof of this result consist an explicit definition of the 2-limit and 2-colimit. We are going to recall these definitions, but morphisms of these categories won't be given in a classical way. Usually they are given as classes of morphisms between

objects of $\mathfrak{b}(i)$ which satisfy some conditions. Here we are going to express them in term of limit and colimit.

Let $\mathfrak{b} : \mathcal{I} \rightarrow \mathcal{CAT}$ be a 2-functor. Let us give some useful notations. Let $i, i' \in \mathcal{I}$ and $\mathcal{I}_{ii'}$ be the category defined by:

- the objects of $\mathcal{I}_{ii'}$ are :

$$\left\{ (i'', s, s') \mid i'' \in \text{Ob}\mathcal{I}, s \in \text{Hom}_{\mathcal{I}}(i, i''), s' \in \text{Hom}_{\mathcal{I}}(i', i'') \right\}$$

visualized by :

$$\begin{array}{ccc} i & \xrightarrow{s} & i'' \\ & \nearrow s' & \\ i' & & \end{array}$$

- the morphisms of $\mathcal{I}_{ii'}$ from (i''_1, s_1, s'_1) to (i''_2, s_2, s'_2) are :

$$\text{Hom}_{\mathcal{I}_{ii'}}((i''_1, s_1, s'_1), (i''_2, s_2, s'_2)) = \{t \in \text{Hom}_{\mathcal{I}}(i''_1, i''_2) \mid t \circ s_1 = s_2, t \circ s'_1 = s'_2\}$$

visualized by :

$$\begin{array}{ccccc} i & & \xrightarrow{s_2} & & i''_2 \\ & \searrow s_1 & & \nearrow t & \\ & & i''_1 & & \\ & \nearrow s'_1 & & \nwarrow s'_2 & \\ i' & & & & \end{array}$$

Lemma 4. *If the category $\mathcal{I}$ is filtered then the category $\mathcal{I}_{ii'}$ is filtered.*

Proof. The proof is straight-foward. $\square$

Proposition 5. *The category $\mathcal{B}$ defined below is a 2-colimit of $\mathfrak{b}$ satisfying the strong factorization property.*

- Objects of $\mathcal{B}$ are pairs (i, X) where $i \in \mathcal{I}$ and $X \in \mathfrak{b}(i)$,
- let (i, X) and (i', Y) be two objects of $\mathcal{B}$, the morphisms from (i, X) to (i', Y) are the elements of the colimit :

$$\text{Hom}_{\mathcal{B}}(X, Y) = \varinjlim_{(i'', s, s') \in \mathcal{I}_{ii'}} \text{Hom}_{\mathfrak{b}(i'')}(b(s)X, b(s')Y)$$

where if $t \in \text{Hom}_{\mathcal{I}_{ii'}}((i''_1, s_1, s'_1), (i''_2, s_2, s'_2))$, its image in the inductive system is the following morphism :

$$\begin{array}{ccc} \text{Hom}_{\mathfrak{b}(i''_1)}(b(s_1)(X), b(s'_1)(Y)) & \longrightarrow & \text{Hom}_{\mathfrak{b}(i''_2)}(b(s_2)(X), b(s'_2)(Y)) \\ h & \longmapsto & b_{t, s'_1}^{-1} \circ b(t)h \circ b_{t, s_1} \end{array}$$

where b_{t, s_1} is the isomorphism given by the 2-functor $\mathfrak{b}$:

$$b_{t, s_1} : b(t \circ s_1) \xrightarrow{\sim} b(t) \circ b(s_1)$$

Proof. See for example [3]. $\square$

If $\mathcal{I}_{ii'}$ is filtered, the morphisms from (i, X) to (i', Y) are the elements of the quotient :

$$\coprod_{i'' \in \mathcal{I}_{ii'}} \text{Hom}_{\mathbf{b}(i'')}(\mathbf{b}(s)X, \mathbf{b}(s')Y) / \sim$$

Where, given $h_1 \in \text{Hom}_{\mathbf{b}(i'_1)}(\mathbf{b}(s_1)X, \mathbf{b}(s'_1)Y)$ and $h_2 \in \text{Hom}_{\mathbf{b}(i'_2)}(\mathbf{b}(s_2)X, \mathbf{b}(s'_2)Y)$, we write $h_1 \sim h_2$ if and only if there exist $(i_3, s_3, s'_3) \in \mathcal{I}_{ii'}$, $t_{13} \in \text{Hom}((i_1, s_1, s'_1), (i_3, s_3, s'_3))$ and $t_{23} \in \text{Hom}((i_2, s_2, s'_2), (i_3, s_3, s'_3))$ such that :

$$b_{t_{13}, s'_1}^{-1} \circ \mathbf{b}(t_{13})h_1 \circ b_{t_{13}, s_1} = b_{t_{23}, s'_2}^{-1} \circ \mathbf{b}(t_{23})h_2 \circ b_{t_{23}, s_2}$$

Similarly we are going to give an explicit construction of a 2-limit. As before morphisms of this category will be given by a limit. Let $\mathbf{c}$ be a contravariant 2-functor :

$$\mathbf{c} : \mathcal{J}^{op} \longrightarrow \mathcal{CAT}$$

Proposition 6. *The category $\mathcal{C}$ define below, is a 2-limit of $\mathbf{c}$ satisfying the strong factorization property.*

- The objects of $\mathcal{C}$ are pairs (X, ϑ^X) where :
 - $X = \{X_j\}_{j \in \mathcal{J}}$, for $X_j \in \text{Ob}(\mathbf{c}(j))$,
 - $\vartheta^X = \{\vartheta_t^X\}_{t \in \text{Hom}_{\mathcal{J}}}$ where, for $t \in \text{Hom}_{\mathcal{J}}(j, j')$, ϑ_t^X is an isomorphism :

$$\vartheta_t^X : X_j \xrightarrow{\sim} \mathbf{c}(t)X_{j'}$$

satisfying the following conditions :

- A) for any $j \in \mathcal{J}$ we have $\text{Id}_j = c_j(X_j) \circ \vartheta_{\text{Id}_j}^X$, where c_j is the morphism given by the 2-functor $\mathbf{c}$:

$$c_j : \mathbf{c}(\text{Id}_j) \xrightarrow{\sim} \text{Id}_{\mathbf{c}(j)}$$

- B) for any two composable morphisms $t : j \rightarrow j'$ and $t' : j' \rightarrow j''$ the following equation holds :

$$\mathbf{c}(t)(\vartheta_{t'}^X) \circ \vartheta_t^X = c_{t, t'}(X_{j''}) \circ \vartheta_{t \circ t'}^X$$

where $c_{t, t'}$ is the isomorphism given by the 2-functor $\mathbf{c}$:

$$c_{t, t'} : \mathbf{c}(t \circ t') \xrightarrow{\sim} \mathbf{c}(t) \circ \mathbf{c}(t')$$

- Let (X, ϑ^X) and (Y, ϑ^Y) be two objects of $\mathcal{C}$, morphisms from (X, ϑ^X) to (Y, ϑ^Y) are elements of the limit :

$$\text{Hom}_{\mathcal{C}}(X, Y) = \varprojlim_{j \in \mathcal{J}} \text{Hom}_{\mathbf{c}(j)}(X_j, Y_j)$$

where if $t \in \text{Hom}_{\mathcal{J}}(j, j')$, its image in the projective system is the following morphism :

$$\begin{array}{ccc} \text{Hom}_{\mathbf{c}(j)}(X_j, Y_j) & \longrightarrow & \text{Hom}_{\mathbf{c}(j')}(X_{j'}, Y_{j'}) \\ h & \longmapsto & (\vartheta_t^Y)^{-1} \circ \mathbf{c}(t)(h) \circ \vartheta_t^X \end{array}$$

This means that a morphism between two objects is the datum of $\{h_j\}_{j \in \mathcal{J}}$, where $h_j \in \text{Hom}_{\mathbf{c}(j)}(X_j, Y_j)$ satisfies the equality :

$$h_j = (\vartheta_t^Y)^{-1} \circ \mathbf{c}(t)(h_{j'}) \circ \vartheta_t^X$$

for all $t : j' \rightarrow j$ morphisms of $\mathcal{J}$.

3 Interchange of filtered 2-colimits and finite 2-limits

We are going to show that filtered 2-colimits commute with finite 2-limits. Let us give first a precise meaning to this sentence.

Let $\mathcal{I}$ be a filtered category, $\mathcal{J}$ a finite category and $\mathbf{a}$ a 2-functor :

$$\mathbf{a} : \mathcal{I} \times \mathcal{J}^{op} \longrightarrow \mathcal{CAT}$$

Let us denote $2\mathfrak{F}(\mathcal{C})$ the 2-category of 2-functors going from the category $\mathcal{C}$ to the 2-category $\mathcal{CAT}$. We have the following proposition, for a proof see for example [3] :

Proposition 7. *The correspondence :*

$$\begin{array}{ccc} 2\mathfrak{F}(\mathcal{C}) & \longrightarrow & \mathcal{CAT} \\ \mathbf{b} & \longmapsto & 2\varinjlim_{i \in \mathcal{I}} \mathbf{b}(i) \end{array}$$

can be extended to a 2-functor between 2-categories. A similar statement holds for 2-limits.

Now let us consider, the natural 2-functor :

$$\begin{array}{ccc} \mathcal{J} & \longrightarrow & 2\mathfrak{F}(\mathcal{I}) \\ j & \longmapsto & \mathbf{a}(\cdot, j). \end{array}$$

The composition of this 2-functor and the one defined in the proposition gives a functor from $\mathcal{J}$ to $\mathcal{CAT}$. As $\mathcal{CAT}$ is complete we can consider its limit. Let us denote

$$2\varprojlim_{j \in \mathcal{J}} 2\varinjlim_{i \in \mathcal{I}} \mathbf{a}(i, j)$$

this limit. We define in the same way the 2-colimit $2\varinjlim_{i \in \mathcal{I}} 2\varprojlim_{j \in \mathcal{J}} \mathbf{a}(i, j)$.

Remark

The composition, for all $i \in \mathcal{I}$ and $j \in \mathcal{J}$, of the functor define by the 2-colimit and the 2-limit :

$$i 2\varprojlim_{j' \in \mathcal{J}} \mathbf{a}(i, j') \longrightarrow \mathbf{a}(i, j) \longrightarrow 2\varinjlim_{i' \in \mathcal{I}} \mathbf{a}(i', j) i$$

defines a functor :

$$\Psi : 2\varinjlim_{i \in \mathcal{I}} 2\varprojlim_{j \in \mathcal{J}} \mathbf{a}(i, j) \longrightarrow 2\varprojlim_{j \in \mathcal{J}} 2\varinjlim_{i \in \mathcal{I}} \mathbf{a}(i, j)$$

Theoreme 8. *The natural functor*

$$\Psi : 2\varinjlim_{i \in \mathcal{I}} 2\varprojlim_{j \in \mathcal{J}} \mathbf{a}(i, j) \longrightarrow 2\varprojlim_{j \in \mathcal{J}} 2\varinjlim_{i \in \mathcal{I}} \mathbf{a}(i, j)$$

is an equivalence of categories.

Proof. The fact that Ψ is fully faithful comes directly from the expression of the morphisms of these two categories in terms of limits and colimits. In detail, using the two propositions 5 and 6, we can give an explicit construction of the categories $2\varinjlim_{i \in \mathcal{I}} 2\varprojlim_{j \in \mathcal{J}} \mathbf{a}(i, j)$ and $2\varprojlim_{j \in \mathcal{J}} 2\varinjlim_{i \in \mathcal{I}} \mathbf{a}(i, j)$.

The category $\varinjlim_{i \in \mathcal{I}} \varprojlim_{j \in \mathcal{J}} \mathbf{a}(i, j)$ is defined as follows :

- its objects are the triples (i, X, ϑ^X) where
 - i is an object of $\mathcal{I}$,
 - $X = \{X_j\}_{j \in \mathcal{J}}$ for X_j is an object of $\mathbf{a}(i, j)$
 - $\vartheta^X = \{\vartheta_t^X\}_{t \in \text{Hom } \mathcal{J}}$, where, for $t : j' \rightarrow j$ an isomorphism of $\mathcal{J}$, ϑ_t^X is a morphism :
$$\vartheta_t^X : X_j \longrightarrow \mathbf{a}(Id_i, t)X_{j'}$$

verifying the two following conditions :

- for any $j \in \text{Obj } \mathcal{J}$ we have $a_{ij}(X_{ij}) \circ \vartheta_{Id_i}^X = Id_{X_i}$, where a_{ij} is the isomorphism $a_{ij} : \mathbf{a}(Id_i, Id_j) \xrightarrow{\sim} Id_{\mathbf{a}(i, j)}$ given by the 2-functor $\mathbf{a}$,
- for any two composable morphisms $t : j \rightarrow j'$ and $t' : j' \rightarrow j''$ the following equation holds :

$$\mathbf{a}(Id_i, t)(\vartheta_{t'}^X) \circ \vartheta_t^X = a_{(Id_i, t), (Id_i, t')}(X_{ij''}) \circ \vartheta_{t' \circ t}$$

where $a_{(Id_i, t), (Id_i, t')} : \mathbf{a}(Id_i, t' \circ t) \xrightarrow{\sim} \mathbf{a}(Id_i, t')\mathbf{a}(Id_i, t)$ is the morphism given by the 2-functor $\mathbf{a}$.

- the set of morphisms from (i, X, ϑ^X) to (i', Y, ϑ^Y) is given by the limit :

$$\text{Hom}(X, Y) := \varinjlim_{i'' \in \mathcal{I}_{i, i'}} \varprojlim_{j \in \mathcal{J}} \text{Hom}_{\mathbf{a}(i'', j)}(\mathbf{a}(s, Id_j)X_j, \mathbf{a}(s', Id_j)Y_j).$$

The category $2\varprojlim_{j \in \mathcal{J}} 2\varinjlim_{i \in \mathcal{I}} \mathbf{a}(i, j)$ is defined as follows :

- the objects are pairs $(\mathcal{X}, \theta^{\mathcal{X}})$, where
 - $\mathcal{X} = \{(i_j, X_{i_j})\}_{j \in \mathcal{J}}$ with $i_j \in \mathcal{I}$ and $X_{i_j} \in \mathbf{a}(i_j, j)$,
 - $\theta^{\mathcal{X}} = \{[\vartheta_t]\}_{t \in \text{Hom } \mathcal{J}}$ where, for $t \in \text{Hom } \mathcal{J}(j, j')$, $[\vartheta_t^{\mathcal{X}}]$ belongs to the quotient :

$$\coprod_{i'' \in \mathcal{I}_{i_j, i_{j'}}} \text{Hom}_{\mathbf{a}(i'', j)}(\mathbf{a}(s, Id_j)X_{i_j}, \mathbf{a}(s', t)X_{i_j}) / \sim$$

and where $[\vartheta_t^{\mathcal{X}}]$ satisfies the following equalities :

$$[a_{ij}(X_{ij}) \circ \vartheta_{Id_i}^X] = [Id_{X_i}] \quad (1)$$

$$[\mathbf{a}(Id_i, t)(\vartheta_{t'}^X) \circ \vartheta_t^X] = [a_{(Id_i, t), (Id_i, t')}(X_{ij''}) \circ \vartheta_{t' \circ t}] \quad (2)$$

and for any two composable morphisms $t : j \rightarrow j'$ and $t' : j' \rightarrow j''$.

- the set of morphisms from $(\mathcal{X}, \theta^{\mathcal{X}})$ to $(\mathcal{Y}, \theta^{\mathcal{Y}})$ is given by the limit :

$$\varprojlim_{j \in \mathcal{J}} \varinjlim_{i'' \in \mathcal{I}_{ij i'_j}} \text{Hom}_{\mathbf{a}(i'', j)}(\mathbf{a}(s, Id_j)X_j, \mathbf{a}(s', Id_j)Y_j)$$

The natural functor between the 2-limits is :

$$\begin{aligned} \varepsilon : \quad 2 \varprojlim_{i \in \mathcal{I}} 2 \varinjlim_{j \in \mathcal{J}} \mathbf{a}(i, j) &\longrightarrow 2 \varprojlim_{j \in \mathcal{J}} 2 \varinjlim_{i \in \mathcal{I}} \mathbf{a}(i, j) \\ (i, \{X_j\}, \{\vartheta_t^{\mathcal{X}}\}) &\longmapsto (\{i, X_j\}, \{[\vartheta_t^{\mathcal{X}}]\}) \end{aligned}$$

Moreover, if $\mathcal{X} = (i, X, \vartheta^{\mathcal{X}})$ and $\mathcal{Y} = (i', Y, \vartheta^{\mathcal{Y}})$ are two objects of $2 \varprojlim_{i \in \mathcal{I}} 2 \varinjlim_{j \in \mathcal{J}} \mathbf{a}(i, j)$,

the morphism from $\text{Hom}_{2 \varprojlim_{i \in \mathcal{I}} 2 \varinjlim_{j \in \mathcal{J}}}(\mathcal{X}, \mathcal{Y})$ to $\text{Hom}_{2 \varprojlim_{j \in \mathcal{J}} 2 \varinjlim_{i \in \mathcal{I}}}(\varepsilon(\mathcal{X}), \varepsilon(\mathcal{Y}))$ induced by ε is the natural morphism :

$$\varprojlim_{i \in \mathcal{I}} \varinjlim_{j \in \mathcal{J}} \text{Hom}(\mathbf{a}(s, Id_j)X_j, \mathbf{a}(s', Id_j)Y_j) \longrightarrow \varprojlim_{j \in \mathcal{J}} \varinjlim_{i \in \mathcal{I}} \text{Hom}(\mathbf{a}(s, Id_j)X_j, \mathbf{a}(s', Id_j)Y_j)$$

As filtered colimits commute with finite limits, the morphism above is an isomorphism and ε is fully faithful.

The proof that Ψ is essentially surjective is similar to the proof of the commutation of filtered limits and finite colimits.

Let $(\{i_j, X_j\}, \{[\theta_t^{\mathcal{X}}]\})$ an object of $2 \varprojlim_{j \in \mathcal{J}} 2 \varinjlim_{i \in \mathcal{I}} \mathbf{a}(i, j)$.

Using the property (ii) of a filtered category inductively, one proves that there exist an object $k' \in \mathcal{I}$ and, for any $j \in \mathcal{J}$, a morphism $s'_j : i_j \rightarrow k'$ in $\mathcal{I}$. Thus, for all t morphism of $\mathcal{J}$, $[\theta_t^{\mathcal{X}}]$ can be viewed as the class of an object $\vartheta_t^{\mathcal{X}}$ of $\text{Hom}(\mathbf{a}(s_j, Id_j)X_{i_j}, \mathbf{a}(s'_j, t)X_{j'})$. Remark that even if the class $[\theta_t^{\mathcal{X}}]$ satisfies the equalities (1) and (2), the objects $\vartheta_t^{\mathcal{X}}$ may not satisfy then. Using the property (iii) of a filtered category inductively, one proves that there exist an object k of $\mathcal{I}$ and a morphism $s_k : k' \rightarrow k$ such that all the equalities hold also for the objects $\vartheta_t^{\mathcal{X}}$. Hence the triple :

$$\left(k, \{ \mathbf{a}(s_k \circ s_j, Id_j)X_{i_j} \}, \{ a_{s_k, s_j}^{-1} \circ \vartheta_t^{\mathcal{X}} \circ a_{s_k, s_{j'}} \} \right)$$

is an object of $2 \varprojlim_{i \in \mathcal{I}} 2 \varinjlim_{j \in \mathcal{J}} \mathbf{a}(i, j)$. For $j \in \mathcal{J}$, the objects X_{i_j} and $\mathbf{a}(s_k \circ s_j, Id_j)X_{i_j}$

are isomorphic and, for all morphism t of $\mathcal{J}$, we have

$$[\vartheta_t^{\mathcal{X}}] = [\theta_t^{\mathcal{X}}]$$

Hence we have shown that Ψ is an equivalence of categories. □

References

- [1] S. Mac Lane. *Categories for the working mathematician. Second edition.* Graduate Texts in Mathematics, 5. Springer-Verlag, 1998.
- [2] R. Street. Categorical structures. *Handbook of algebra*, 1996.
- [3] I. Waschies. The stack of microlocal perverse sheaves. *Bull. Soc. Math. France* 132, 2004.